

Quenched Charmed Meson Spectra using Tadpole Improved Quark Action on Anisotropic Lattices¹

Liuming Liu^a, Shiquan Su^a, Xin Li^a and Chuan Liu^a

^a*School of Physics
Peking University
Beijing, 100871, P. R. China*

Abstract

Charmed meson charmonium spectra are studied with improved quark actions on anisotropic lattices. We measured the pseudo-scalar and vector meson dispersion relations for 4 lowest lattice momentum modes with quark mass values ranging from the strange quark to charm quark with 3 different values of gauge coupling β and 4 different values of bare speed of light ν . With the bare speed of light parameter ν tuned in a mass-dependent way, we study the mass spectra of D , D_s , η_c , D^* , D_s^* and J/ψ mesons. The results extrapolated to the continuum limit are compared with the experiment and qualitative agreement is found.

Key words: Non-perturbative renormalization, improved actions, anisotropic lattice, charmed meson spectrum.

PACS: 12.38.Gc, 11.15.Ha

1 Introduction

In recent years, anisotropic lattices and improved lattice actions have been used extensively in lattice QCD calculations. In this letter, we present our preliminary numerical study on the spectrum of charmed mesons and charmoniums. The gauge action employed in this work is the tadpole improved

¹ This work is supported by the Key Project of National Natural Science Foundation of China (NSFC) under grant No. 10421003, No. 10235040 and supported by the Trans-century fund from Chinese Ministry of Education.

gluonic action on asymmetric lattices as in Ref. [2]: Using such a gauge action, glueball and light hadron spectrum has been studied within the quenched approximation [1,2,3,4,5,6,7].

It has been suggested that relativistic heavy quarks can also be treated with the help of anisotropic lattices (the Fermi lab approach), with improvements [8,9,10,11,12]. Using various versions of the quark actions, charmed meson spectrum, charmonium spectrum, charmed baryons have been studied on the lattice [13,14,15,16,17,18]. However, in order to take full advantage of the improved quark action on anisotropic lattices, some parameters in the action have to be tuned, either perturbatively or non-perturbatively, in order to gain as much improvement as possible. Some numerical studies of these parameters have already appeared in the literature [19,12,20,21,22]. In particular, it turns out that the bare speed of light parameter ν has to be tuned non-perturbatively in a quark *mass-dependent* way as is shown in Ref. [23]. With the parameter ν tuned as such, we can study quenched meson spectrum with quark mass values ranging from the strange to the charm, using the same quark action.

The anisotropic quark actions used in our study is: $S_f = \sum_{xy} \bar{\psi}_x \mathcal{M}_{xy} \psi_y$ with the fermion matrix given by [23]:

$$\begin{aligned} \mathcal{A}_{xy} = & \delta_{xy} \left[1/(2\kappa_{max}) + \rho_t \sum_{i=1}^3 \sigma_{0i} \mathcal{F}_{0i} + \rho_s (\sigma_{12} \mathcal{F}_{12} + \sigma_{23} \mathcal{F}_{23} + \sigma_{31} \mathcal{F}_{31}) \right] \\ & - \sum_{\mu} \eta_{\mu} \left[(1 - \gamma_{\mu}) U_{\mu}(x) \delta_{x+\mu,y} + (1 + \gamma_{\mu}) U_{\mu}^{\dagger}(x - \mu) \delta_{x-\mu,y} \right] . \end{aligned} \quad (1)$$

The coefficients in the matrix are given explicitly by:

$$\begin{aligned} \eta_i = \frac{\nu}{2u_s} , \quad \eta_0 = \frac{\xi}{2} , \quad \sigma = \frac{1}{2\kappa} - \frac{1}{2\kappa_{max}} , \\ \rho_t = \nu \frac{(1 + \xi)}{4u_s^2} , \quad \rho_s = \frac{\nu}{2u_s^4} . \end{aligned} \quad (2)$$

In this letter, we present our preliminary quenched results for the meson spectra using the above improved quark action on anisotropic lattices.

2 Simulation Results

The whole procedure of the calculation involves three major steps: In the first step, we determine the optimal value of the bare speed of light parameter ν , using energy-momentum dispersion relations of pseudo-scalar mesons; In the

second step, the masses of the mesons that we are interested in are extrapolated/interpolated to the optimal value of ν ; In the third step, we perform chiral and continuum extrapolations as usual.

The lattices in this study are of size $6 \times 9 \times 12 \times 50$ for $\beta = 2.4, 2.6$ and $8 \times 12 \times 16 \times 50$ for $\beta = 2.8$. The spatial lattice spacing a_s roughly range from $0.13fm$ and $0.23fm$ in physical units. The aspect ratio is $\xi = a_s/a_t \simeq \xi_0 = 5$ for all lattices. For gauge field configurations at a given value of β , 4 different values of the bare speed of light parameter ν and 12 different values of the hopping parameter κ are studied. The range of κ roughly covers the quark mass from the strange to the charm. About 300 configurations are used for each parameter set in this study.

The energy values of meson with definite three-momentum $\mathbf{k}$ (including zero-momentum) is obtained from their respective correlation functions by finding the plateaus in their effective mass plots. The errors for the data points in the effective mass plots are analyzed using the standard jack-knife method. We obtain from the pseudo-scalar and vector channel the corresponding energy $E_{PS/V}(m_1, m_2, \nu_1, \nu_2, \mathbf{k})$, where m_1 and m_2 are the bare quark mass parameters defined by: $m \equiv 1/(2\kappa) - 1/[2\kappa_{cr}(\nu)]$. The value of $1/[2\kappa_{cr}(\nu)]$ is obtained by the condition of vanishing pseudo-scalar meson mass for each ν . Although for different values of ν , the set of values for m turns out to be slightly different, we always interpolate/extrapolate them to the same set of values [23].

To obtain the optimal value of ν , the values of $E_{PS}^2(m, m, \nu, \nu, \mathbf{k})$ are fitted linearly against the three-momentum squared $\mathbf{k}^2$ in the low-momentum region:

$$E_{PS}^2(m, m, \nu, \nu, \mathbf{k}) = M_{PS}^2(m, m, \nu, \nu) + Z_{PS}(m, m, \nu, \nu) \mathbf{k}^2 . \quad (3)$$

The condition: $Z_{PS}[m, m, \nu_{opt}(m), \nu_{opt}(m)] = 1$ then yields the optimal value of ν as a function of the bare quark mass parameter: $\nu_{opt}(m)$. In practice, we use the values of $Z_{PS}(m, m, \nu, \nu)$ at different ν to perform a linear extrapolation/interpolation in ν for each value of m .

We now proceed to the second step of our study, namely to extrapolate/interpolate the meson mass values to the optimal values of ν obtained from the pseudo-scalar dispersion relations. This is achieved by a linear extrapolation/interpolation in ν , or equivalently in the quantity $Z_{PS}(m, m, \nu, \nu)$.

Similar procedure discussed above can be repeated for mesons consisting of a quark and an anti-quark with different mass. We interpolate/extrapolate the meson masses to the optimal bare speed of light parameter. The outcome of this procedure are the quantities of $M_{PS/V}^2(m_1, m_2, \nu_{opt}(m_1), \nu_{opt}(m_2))$. Since these quantities only depend on the bare quark mass parameter pairs (m_1, m_2) , we denote them by $\bar{M}_{PS/V}^2(m_1, m_2)$.

In the third step of the analysis, we perform the chiral extrapolations. First, we determine the physical strange and charm bare quark mass parameters: $m_s^{(phy)}$ and $m_c^{(phy)}$. In this work, we use the mass of the vector meson with the same flavor of quarks to determine $m_s^{(phy)}$ and $m_c^{(phy)}$. The results of $M_V^2(m, m, \nu_{opt}(m), \nu_{opt}(m))$ are fitted quadratically versus the quark mass parameter:

$$a_t^2 M_V^2[m, m, \nu_{opt}(m), \nu_{opt}(m)] = A + Bm + Cm^2. \quad (4)$$

We perform two such quadratic fits, one in the low quark mass region, the other in heavy quark mass region. The scale of the lattice is set by $r_0 = 0.5\text{fm}$ (the so-called Sommer scale). Then the value of $m_s^{(phy)}$ and $m_c^{(phy)}$ are obtained by requiring the vector meson mass obtained from the simulation to exactly reproduce the mass of the physical Φ and the physical J/ψ meson mass. Fig. 1 shows the quadratic extrapolations of $M_V^2(m, m, \nu_{opt}(m), \nu_{opt}(m))$ versus the bare quark mass m at $\beta = 2.6$. The physical Φ and J/ψ mass are also lined out by horizontal lines in the figure. The intersection point of the lower/higher horizontal line representing $a_t^2 M_\Phi^2/a_t^2 M_{J/\psi}^2$ and the quadratic fitting curve yields the estimate for the physical strange quark mass parameter $m_s^{(phy)}$ and $m_c^{(phy)}$, respectively.

Having determined the values of $m_s^{(phy)}$ and $m_c^{(phy)}$, we then make quadratic extrapolations/interpolations for all the meson mass squared ($\bar{M}_{P/S/V}^2(m_1, m_2)$) versus the quark mass parameters and extrapolate/interpolate them to their physical values $m_{s/c}^{(phy)}$, depending on which type of meson is being considered. The final results are the mass values of various charmed mesons and charmonium states at different β . These results are listed in Table tab:spectra. ²

Table 1

Extracted physical mass (in unit of MeV) of vector and pseudo scalar meson made up of one charm quark and another quark, for different β and continuum limit. For comparison, we also list the experimental values for the mesons in the last column.

β	2.8	2.6	2.4	continuum limit	Experiment
$m(D)$	1835(35)	1862(5)	1860(9)	1847(68)	1869
$m(D_s)$	1976(21)	1993(3)	1959(5)	2164(39)	1969
$m(\eta_c)$	2992(2)	3037(1)	3020(10)	2843(23)	2980
$m(D^*)$	2015(27)	1965(6)	1993(33)	2046(131)	2007
$m(D_s^*)$	2140(18)	2087(3)	2080(13)	2192(67)	2112

Finally, the results of the meson masses in the continuum are obtained by extrapolate their values at finite lattice spacings to the continuum limit. In Fig. 2, we show the mass ratios of various mesons over the mass of the J/ψ

² The mass of J/ψ (not included in Table 1) is by definition that of the experimental value since we have used the mass of J/ψ to determine the bare charm quark mass.




Fig. 1. Quadratic extrapolation/interpolation of $M_V^2(m, m, \nu_{opt}(m), \nu_{opt}(m))$ versus the bare quark mass m at $\beta = 2.6$. The two curves are the fits in the light and heavy quark mass regions, respectively. The corresponding fitting qualities and ranges are also shown in the figure. The fitting ranges are self-adjusted to yield minimum χ^2 per degree of freedom. The horizontal and vertical lines indicate the value for the physical Φ and J/ψ meson and the fixed value of bare quark mass $m_c^{(phy)}$ and $m_c^{(phy)}$, respectively.

versus lattice spacing a_s . Linear fits are performed for these mass ratios and the results are indicated by the straight lines. The extrapolated results are also indicated by the points at $a_s = 0$. The numerical results are also listed in Table 1. The fitting quality for most cases are reasonable. However, since we have only 3 different values of β for this preliminary study, the continuum limit extrapolations suffer big errors. Needless to say, more points at different values of a_s , preferably at smaller a_s , will help to improve the final result at the continuum limit.

To compare with the experiment, we have also tabulated the results from Particle Data Group (PDG) in the last column of Table 1. It is seen that the qualitative agreement is good, especially for D , D^* and D_s^* mesons. The mass values of the D_s and η_c are away from the experimental value by 5σ .




Fig. 2. Continuum limit extrapolation for each meson mass over J/ψ meson mass. Each panel indicates different meson. Data points (shown by green points) in the plot are ratios of the mass of each meson to the mass of J/ψ for different β (corresponding to different value of a_s/r_0). The red points represent the physical mass ratios at the continuum limit.

These disagreements might come from the fact that our lattice spacing is still somewhat large and the continuum extrapolation is not very well controlled. Of course, quenching effects are also present which might be important for the fine structure of the spectrum.

3 conclusion

In this paper, we present a preliminary study of the quenched meson spectra for the charmed mesons and charmonium states. To perform such a calculation using the tadpole improved anisotropic Wilson quark action, we tune the bare speed of light parameter ν in a quark mass dependent way with quark mass values ranging from the strange to the charm. The optimal values of ν are obtained for various values of β using the pseudo-scalar meson dispersion

relations. Using the tadpole improved anisotropic Wilson action with these optimized parameters, we determine the ground state meson spectra for D , D_s , η_c , D^* , D_s^* and J/ψ mesons. The results are extrapolated towards the continuum limit where qualitative agreement with the experiment is found. With the method outlined in this preliminary study, we plan to carry out spectrum calculations of other excited meson states, baryon states and possibly hybrid states involving a charm quark in the future.

References

- [1] C. Morningstar and M. Peardon. *Phys. Rev. D*, 56:4043, 1997.
- [2] C. Morningstar and M. Peardon. *Phys. Rev. D*, 60:034509, 1999.
- [3] C. Liu. *Chinese Physics Letter*, 18:187, 2001.
- [4] C. Liu. *Communications in Theoretical Physics*, 35:288, 2001.
- [5] C. Liu. In *Proceedings of International Workshop on Nonperturbative Methods and Lattice QCD*, page 57. World Scientific, Singapore, 2001.
- [6] C. Liu and J. P. Ma. In *Proceedings of International Workshop on Nonperturbative Methods and Lattice QCD*, page 65. World Scientific, Singapore, 2001.
- [7] C. Liu. *Nucl. Phys. (Proc. Suppl.) B*, 94:255, 2001.
- [8] A.X. El-Khadra, A.S. Kronfeld, and P.B. Mackenzie. *Phys. Rev. D*, 55:3933, 1997.
- [9] M. Alford, T. R. Klassen, and P. Lepage. *Phys. Rev. D*, 58:034503, 1998.
- [10] T. R. Klassen. *Nucl. Phys. (Proc. Suppl.) B*, 73:918, 1999.
- [11] J. Harada, A.S. Kronfeld, H. Matsufuru, N. Nakajima, and T. Onogi. *Phys. Rev. D*, 64:074501, 2001.
- [12] J. Harada, H. Matsufuru, T. Onogi, and A. Sugita. *Phys. Rev. D*, 66:014509, 2002.
- [13] P.B. Mackenzie, S.M. Ryan, and J.N. Simone. *Nucl. Phys. (Proc. Suppl.) B*, 63:305, 1998.
- [14] P. Chen. *Phys. Rev. D*, 64:034509, 2001.
- [15] R. Lewis, N. Mathur, and R.M. Woloshyn. *Phys. Rev. D*, 64:094509, 2001.
- [16] M. Okamoto et al. *Phys. Rev. D*, 65:094508, 2002.
- [17] A. Juettner and J. Rolf. *Phys. Lett. B*, 560:59, 2003.

- [18] Y. Nemoto, N. Nakajima, H. Matsufuru, and H. Suganuma. *Phys. Rev. D*, 68:094505, 2003.
- [19] Junhua Zhang and C. Liu. *Mod. Phys. Lett. A*, 16:1841, 2001.
- [20] T. Umeda et al. *Phys. Rev. D*, 68:034503, 2003.
- [21] S. Hashimoto and M. Okamoto. *Phys. Rev. D*, 67:114503, 2003.
- [22] Justin Foley, Alan O Cais, Mike Peardon, and Sinead M. Ryan. *hep-lat/0405030*, 2004.
- [23] Shi-Quan Su, Liu-Ming Liu, and Chuan Liu. *hep-lat/0412034*, accepted by Int. J. Mod. Phys. A, 2005.